

Advice on the use of masks for children in the community in the context of COVID-19

Annex to the Advice on the use of masks in the context of COVID-19

21 August 2020



Purpose of the document

This document provides guidance to decision makers, public and child health professionals to inform policy on the use of masks for children in the context of the COVID-19 pandemic. It does not address the use of masks for adults working with children or parents/guardians or the use of masks for children in health-care settings. This interim guidance will be revised and updated as new evidence emerges.

Background

The World Health Organization (WHO) and the United Nations Children's Fund (UNICEF) advise the use of masks according to a risk-based approach, as part of a comprehensive package of public health interventions that can prevent and control the transmission of certain viral respiratory diseases, including COVID-19. Compliance with other measures including physical distancing, hand hygiene, respiratory etiquette and adequate ventilation in indoor settings is essential for reducing the spread of SARS-CoV-2, the virus that causes COVID-19.

This guidance provides specific considerations for the use of non-medical masks, also known as fabric masks, by children as a means for source control in the context of the current COVID-19 pandemic. The document is an annex to the WHO's Advice on the use of masks in the context of COVID-19¹ in which further details on fabric masks can be found. This annex also advises the use of medical masks for children under certain conditions. For the purposes of this guidance, children are defined as anyone below the age of 18 years².

Methodology for developing the guidance

The World Health Organization (WHO) Infection Prevention and Control (IPC) Guidance Development Group (GDG) and experts from UNICEF and the International Paediatric Association (IPA) jointly reviewed the available evidence to develop guidance on the use of masks for children in the context of the current pandemic. Five international expert meetings were held between June and August 2020. In the absence of strong scientific evidence, consensus among these groups forms the main basis for this guidance. Furthermore, the draft guidance was reviewed by a multidisciplinary group of additional external experts prior to finalization.

Available evidence

Transmission of COVID-19 in children

Currently, the extent to which children contribute to transmission of SARS-CoV-2 is not completely understood. According to the WHO global surveillance database of laboratory-confirmed cases developed from case report forms provided to WHO by Member States³ and other studies, 1-7% of COVID-19 cases are reported to be among children, with relatively few deaths compared to other age groups⁴⁻⁸. The European Centre for Disease Prevention and Control (ECDC) has recently reported the age distribution of COVID-19 among children in the European Union (EU), European Economic Area (EEA) and the United Kingdom (UK); they reported that as of 26 July 2020, 4% of all cases in the EU/EEA and the UK were among children⁶.

To date, the available evidence suggests that most reported cases among children have resulted from transmission within households, although this observation may have been influenced by school closures and other stay at home measures implemented by some countries^{7,9}. Although culture-competent virus has been isolated from symptomatic children with viral load levels found to be similar to that in adults¹⁰, evidence from available studies of contacts of COVID-19 cases and cluster investigations suggests that children are unlikely to be the main drivers of COVID-19 transmission^{7,9 11-14}. To date, documented transmission among children and staff within educational settings is limited¹⁵⁻²⁰. Evidence is also limited regarding the prevalence of SARS-CoV-2 infection among children, as measured by seroepidemiology studies. However, available evidence suggests that seroprevalence appears to be lower for younger children compared to older children and adults^{17,21-25}.

Studies of viral load and the duration of viral shedding of infectious virus in children compared to adults, are also limited. One published study suggests that viral load in infected patients may differ by age, and that symptomatic children have a longer duration of viral shedding than asymptomatic children²⁵. Some studies have reported that children below five years are reported to have lower

amounts of viral RNA in respiratory secretions and faeces compared to school children, adolescents and adults^{26,27}. However, one study from the United States of America found that children below five years with mild to moderate COVID-19 have higher amounts of viral RNA in their upper respiratory samples compared with older children and adults²⁸, while a pre-print (non-peer-reviewed) study from Germany reported no differences in the amount of viral RNA among adults and children²⁹.

In summary, the degree to which age alone, regardless of symptoms, affects viral load and transmission is not well understood.

Available evidence on the use of masks in children for COVID-19 and other respiratory diseases

Evidence on the benefits and harms of children wearing masks to mitigate transmission of COVID-19 and other coronaviruses is limited. However, some studies have evaluated the effectiveness of mask use in children for influenza and other respiratory viruses³⁰⁻³⁴. A study of mask wearing during seasonal influenza outbreaks in Japan noted that the use of masks was more effective in higher school grades (9-12 year old children in grades 4-6) than lower grades (6-9 year old children, in grades 1-3)³⁴. One study, conducted under laboratory conditions and using non-betacoronaviruses, suggested that children between five and 11 years old were significantly less protected by mask wearing compared to adults, possibly related to inferior fit of the mask³⁵. Other studies found evidence of some protective effect for influenza for both source control³⁰ and protection in children³⁴, although overall compliance with consistent mask wearing, especially among children under the age of 15, was poor.

Some studies, including studies conducted in the context of influenza and air pollution, found the use and acceptability of mask wearing to be highly variable among children, ranging from very low to acceptable levels and decreasing over time while wearing masks^{30,31,33,36-38}. One study was carried out among primary school children during COVID-19 and reported 51.6% compliance.³¹

Several studies found that factors such as warmth, irritation, breathing difficulties, discomfort, distraction, low social acceptability and poor mask fit were reported by children when using masks^{30,33,36,37}. So far, the effectiveness and impact of masks for children during play and physical activity have not been studied; however, a study in adults found that N95 respirator and surgical masks reduced cardiopulmonary capacity during heavy exertion³⁹.

Main conclusions

According to the limited available evidence, young children may have lower susceptibility to infection compared to adults^{11,14}, however available data suggests that this may vary by age among children^{17, 21-25}. Data from seroepidemiology studies and transmission studies suggest that older children (e.g. teenagers) may play a more active role in transmission than younger children.^{11,14,17, 21-25}

The benefits of wearing masks in children for COVID-19 control should be weighed against potential harm associated with wearing masks, including feasibility and discomfort, as well as social and communication concerns. Factors to consider also include age groups, sociocultural and contextual considerations and availability of adult supervision and other resources to prevent transmission.

There is a need for data from high quality prospective studies in different settings on the role of children and adolescents in transmission of SARS-CoV-2⁴⁰, on ways to improve acceptance and compliance of mask use and on the effectiveness of masks use in children. These studies must be prioritized and include prospective studies of transmission within educational settings and households stratified by age groups (ideally <2, 2-4, 5-11 and > 12 years) and with different prevalence and transmission patterns. Particular emphasis must be placed on studies in schools in low- and middle-income settings.

Advice to decision makers on the use of masks for children in the community

Overarching guiding principles

Given the limited evidence on the use of masks in children for COVID-19 or other respiratory diseases, including limited evidence about transmission of SARS-CoV-2 in children at specific ages, the formulation of policies by national authorities should be guided by the following overarching public health and social principles:

- Do no harm: the best interest, health and well-being of the child should be prioritized.
- The guidance should not negatively impact development and learning outcomes.
- The guidance should consider the feasibility of implementing recommendations in different social, cultural and geographic contexts, including settings with limited resources, humanitarian settings and among children with disabilities or specific health conditions.

Advice on the use of masks in children

WHO and UNICEF advise decision makers to apply the following criteria for use of masks in children when developing national policies, in countries or areas where there is known or suspected community transmission^a of SARS-CoV-2 and in settings where physical distancing cannot be achieved.

1. Based on the expert opinion gathered through online meetings and consultative processes, children aged up to five years should not wear masks for source control. This advice is motivated by a “do no harm” approach and considers:
 - childhood developmental milestones^{b 41}
 - compliance challenges and
 - autonomy required to use a mask properly.

The experts (following the methods described above) recognized that the evidence supporting the choice of the age cut-off is limited (see above, section related to transmission of COVID-19 in children), and they reached this decision mainly by consensus. The rationale included consideration of the fact that by the age of five years, children usually achieve significant developmental milestones, including the manual dexterity and fine motor coordination movements needed to appropriately use a mask with minimal assistance.

In some countries, guidance and policies recommend a different and lower age cut-off for mask use⁴²⁻⁴⁵. It is recognized that children may reach developmental milestones at different ages and children five years of age and under may have the dexterity needed to manage a mask. Based on the do no harm approach, if the lower age cut-off of two or three years of age is to be used for recommending mask use for children, appropriate and consistent supervision, including direct line of sight supervision by a competent adult and compliance need to be ensured, especially if mask wearing is expected for an extended period of time. This is both to ensure correct use of the mask and to prevent any potential harm associated with mask wearing to the child.

Children with severe cognitive or respiratory impairments who have difficulties tolerating a mask should, under no circumstances, be required to wear masks.

Other IPC, public health and social measures should be prioritized to minimize the risk of SARS-CoV-2 transmission for children five years of age and under; specifically maintaining physical distance of at least 1 meter where feasible, educating children to perform frequent hand hygiene and limiting the size of school classes. It is also noted that there may be other specific considerations, such as the presence of vulnerable persons or other local medical and public health advice that should be considered when determining if children five years of age and under need to wear a mask.

2. For children between six and 11 years of age, a risk-based approach should be applied to the decision to use of a mask. This approach should take into consideration:
 - intensity of transmission in the area where the child is and updated data/available evidence on the risk of infection and transmission in this age group;
 - social and cultural environment such as beliefs, customs, behaviour or social norms that influence the community and population’s social interactions, especially with and among children;
 - the child’s capacity to comply with the appropriate use of masks and availability of appropriate adult supervision;
 - potential impact of mask wearing on learning and psychosocial development; and
 - additional specific considerations and adaptations for specific settings such as households with elderly relatives, schools, during sport activities or for children with disabilities or with underlying diseases.
3. Advice on mask use in children and adolescents 12 years or older should follow the WHO guidance for mask use in adults¹ and/or the national mask guidelines for adults.

Even where national guidelines apply, additional specific considerations (see below) and adaptations for special settings such as schools, during sport, or for children with disabilities or with underlying diseases will need to be specified.

4. The use of a medical mask for immunocompromised children or for paediatric patients with cystic fibrosis or certain other diseases (e.g. cancer) is usually recommended but should be assessed in consultation with the child’s medical provider^{46,47}.

^a Defined by WHO as “experiencing larger outbreaks of local transmission defined through an assessment of factors including, but not limited to: large numbers of cases not linkable to transmission chains; large numbers of cases from sentinel surveillance; and/or multiple unrelated clusters in several areas of the country/territory/area” (<https://www.who.int/publications-detail/global-surveillance-for-covid-19-caused-by-human-infection-with-covid-19-virus-interim-guidance>)

^b An example of considering childhood developmental milestones as defined by CDC are available here:

https://www.cdc.gov/ncbddd/actearly/pdf/checklists/Checklists-with-Tips_Reader_508.pdf
https://www.cdc.gov/ncbddd/actearly/pdf/checklists/Checklists-with-Tips_Reader_508.pdf

For children of any age with developmental disorders, disabilities or other specific health conditions that might interfere with mask wearing, the use of masks should not be mandatory and should be assessed on a case by case basis by the child's educator and/or medical provider.

Implementation considerations

Local epidemiology and contextual issues, such as intensity of transmission, ability to physically distance or implement appropriate ventilation measures in indoor settings, age mixing and contact with other vulnerable individuals should be considered when adopting advice for wearing masks among different age groups, in addition to potential harms and adverse effects of mask wearing.

Age-appropriate communication aimed at improving understanding of the purpose of mask wearing, safe and appropriate mask wearing and maintenance of masks, should be provided by parents/guardians, teachers, educators, and trusted community members through role-modelling. Materials, messages and mechanisms for communication on masks for children should remain flexible and adaptive and be systematically reviewed and updated based on changes in evidence and community needs and questions^{48,49}. Children should also be listened to regarding their perceptions and any concerns about wearing a mask. Adapted communication should be available for different social, cultural and linguistic settings, with feedback mechanisms in place for responding to children's questions and expectations.

Specific education and communication messages should be developed to ensure that the use of masks does not result in a false sense of security or disregard for other public health measures by children. It is important to emphasize that the use of masks is one tool and that children should also adhere to physical distancing, hand hygiene and respiratory etiquette. Parents, family members, teachers and educators have a critical role in ensuring that these messages are consistently conveyed to children.

Strategies for assisting children, especially in younger age groups, to manage the wearing of masks safely and effectively should be included in the implementation of this advice. This may include processes for safe storage of used masks for reuse by the same child after eating or exercising, storing soiled masks (e.g. in dedicated bags or containers) before they can be laundered and storage and supply of additional clean masks if a child's mask becomes soiled, wet, or is lost.

Masks should be made accessible free of charge to children living in households or geographic areas with social vulnerabilities and limited resources to ensure equitable access for all children. Consideration should also be made for provision of masks for the journey to and from school.

The design of face masks for children should take into consideration the overall quality of the fabric, suitable breathability and comfort¹ and child-friendliness (appropriate size, colours, design, etc.) to help improve their acceptance of and use by children. Specific attention needs to be given to the care of masks and the need for masks to be changed when they get wet or soiled. Specific measures will need to be in place for children under 12 years who are in a situation where they are asked to wear masks.

The age cut-off for wearing a mask should be adapted to social or school settings to avoid stigmatizing and alienating children in mixed-aged groups where individuals may be on opposite sides of a recommended age cut-off. For example, in situations where older children for whom masks are advised are in the same class as younger children who fall below the age cut-off for wearing masks, the older learners might be exempt from wearing masks.

Specific additional considerations for children with disabilities

Children with developmental disorders or disabilities may face additional barriers, limitations and risks and therefore should be given alternative options to mask wearing, such as face shields (see below). Policies on masks should be adapted for children with disabilities based on social, cultural and environmental considerations.

Some children with disabilities require close physical contact with therapists, educators or social workers. In this context, it is critical that all care providers adopt key IPC measures, including wearing masks, and that settings are adapted to strengthen IPC.

The wearing of masks by children with hearing loss or auditory problems may present learning barriers and further challenges, exacerbated by the need to adhere to the recommended physical distancing⁵⁰. These children may miss learning opportunities because of the degraded speech signal stemming from mask wearing, the elimination of lipreading and speaker expressions and physical distancing. Adapted masks to allow lipreading (e.g. clear masks) or use of face shields (see below) may be explored as an alternative to fabric masks⁵¹.

Specific additional considerations for school settings

To facilitate the operationalization of this guidance in school settings (as per national standards) it is advised that the age categories be adapted to the national/local education level structure.

The use of masks by children and adolescents in schools should only be considered as one part of a comprehensive strategy to limit the spread of COVID-19. The following guidance documents can be used to inform policy making and programming either for a comprehensive school safety strategy when re-opening or operations in the context of COVID-19:

- [WHO considerations for school-related public health measures in the context of COVID-19](#)
- [WB/WFP/UNESCO/UNICEF framework for school reopening](#)
- [WHO/UNICEF/IFRC Interim Guidance for COVID-19 Prevention and Control in Schools](#)

As part of the comprehensive school safety strategy for reopening, the views of teachers and educators on the perception of risks and the time burden required to ensure adherence to COVID-19 policies in schools and classrooms—including the use of masks by children—should be considered. Situations where wearing a mask can significantly interfere with the learning process and have a negative impact on critical school activities like physical education, meal programs, play time and sports – as well as learning – require special consideration.

If wearing of fabric masks is recommended in schools, specific instructions and supplies should be provided for the safe storage, handling and availability of fabric masks (see above). A sufficient supply of appropriate masks should be ensured for all school children. Basic water, sanitation and hygiene requirements should be met in the school building so that comprehensive IPC measures can be implemented, linked to specific age-appropriate educational activities.

If medical or disposable masks are used in specific situations, a system for waste management including disposal of used masks will need to be established to reduce the risk of contaminated masks being disposed of in classrooms and playgrounds.

No children should be denied access to education because of mask wearing or the lack of a mask because of low resources or unavailability⁵⁰.

Alternative to fabric masks for children

Face shields

Face shields are designed to be used⁵² to provide protection from splashes of biological fluid (particularly respiratory secretions), chemical agents and debris^{53,54} into the eyes. In the context of protection from SARS-CoV-2 transmission through respiratory droplets, they are used by health workers as personal protective equipment (PPE) for eye protection in combination with a medical mask or a respirator^{55,56}. In the context of COVID-19 in community settings, some children may not be able to wear a mask for a variety of reasons (e.g. health issues, fear of mask), and thus, face shields may be considered as an alternative to masks as respiratory droplet protection or as source control, based on availability, improved feasibility and better tolerability^{57,58}. Some countries, such as Australia⁵⁹ recommend face shields as an alternative to a mask. Other countries, such as Singapore⁶⁰ advise that both a mask and a shield can be worn together, but acknowledge that children with special needs may need to be exempt from wearing either.

WHO and UNICEF have reviewed the current available evidence on the use of face shields for respiratory droplet protection and/or source control in the context of the COVID-19 pandemic. While a face shield may confer partial protection⁵⁵ of the facial area against respiratory droplets with the added benefit of ease of use, the effectiveness of face shields for source control has not yet been adequately studied. Droplets may be exhaled or inhaled from the open gaps between the visor and the face⁵², which is a disadvantage inherent to its design⁵³. Other design disadvantages include glaring, fogging, optical imperfection, and being bulkier than goggles and safety glasses⁶¹. There are many emerging face shield designs that attempt to overcome these limitations, but current laboratory testing standards only assess face shields for their ability to provide eye protection from chemical splashes^{61,62}. Further research and laboratory challenge standards are urgently needed to investigate the effectiveness of face shields for respiratory droplet protection and/or source control⁵⁶. At present, face shields are considered to provide a level of eye protection only and should not be considered as an equivalent to masks with respect to respiratory droplet protection and/or source control.

WHO and UNICEF will continue to monitor emerging information on the use of face shields for the prevention of respiratory virus transmission. WHO and UNICEF advise that when physical distance cannot be maintained, and in special situations where it is not practical to wear a mask (for example, among children with hearing loss or other disabilities or health conditions that limit compliance with wearing fabric or medical masks and consequently their utility), face shields may be used while taking the following considerations into account:

- The face shield is an incomplete physical barrier and does not provide the filtration layers of a mask.
- The face shield should cover the entire face, be wrapped around the sides of the face and extend to below the chin⁵⁸.
- Reusable face shields must be properly cleaned (with soap or a detergent and water), disinfected (with 70-90% alcohol) and stored after each use⁴⁴. Face shields that will withstand the use of disinfectants without damaging their optical properties should be selected.
- Maintaining physical distance of at least 1 m (3.3 feet) should be maintained where feasible, with ongoing promotion of frequent hand hygiene and respiratory etiquette⁵⁶.
- Caution should be taken to avoid injury when children don, wear, and doff face shields.

Monitoring and evaluation of the impact of the use of masks in children

If authorities decide to recommend mask-wearing for children, key information should be collected on a regular basis to accompany and monitor the intervention. Monitoring and evaluation should be established at the onset and should include indicators that measure the impact on the child's health, including mental health; reduction in transmission of SARS-CoV-2; motivators and barriers to mask wearing; and secondary impacts on a child's development learning, attendance in school, ability to express him/herself or access school; and impact on children with developmental delays, health conditions, disabilities or other vulnerabilities.

Data should be used to inform strategies on communication; training and support to teachers, educators, and parents; engagement activities for children; and distribution of materials that empower children to use masks appropriately.

Analysis should include sex, age, physical, social and economic stratification to ensure that the policy implementation contributes to reducing health and social inequities.

WHO and UNICEF will continue to closely monitor emerging evidence about this topic and the situation for any changes that may affect this interim guidance. Should any factors change, WHO and UNICEF will issue a further update. Otherwise, this interim guidance document will expire six months after the date of publication.

Acknowledgements

This document was developed in consultation with the following members of:

1. The WHO Health Emergencies Programme (WHE) Ad-hoc COVID-19 Guidance Development Group:
Jameela Alsalman, Ministry of Health, Bahrain; Anucha Apisarnthanarak, Thammasat University Hospital, Thailand; Baba Aye, Public Services International, France; Roger Chou, Oregon Health Science University, USA; May Chu, Colorado School of Public Health, USA; John Conly, Alberta Health Services, Canada; Barry Cookson, University College London, United Kingdom (UK); Nizam Damani, Southern Health & Social Care Trust, UK; Dale Fisher, Goarn, Singapore; Tiouiri Benaissa Hanene, CHU La Rabta Tunisia; Joost Hopman, Radboud University Medical Center, The Netherlands; Mushtuq Husain, Institute of Epidemiology, Disease Control & Research, Bangladesh; Kushlani Jayatilleke, Sri Jayewardenapura General Hospital, Sri Lanka; Seto Wing Jong, School of Public Health, Hong Kong SAR, China; Souha Kanj, American University of Beirut Medical Center, Lebanon; Daniele Lantagne, Tufts University, USA; Fernanda Lessa, Centers for Disease Control and Prevention, USA; Anna Levin, University of São Paulo, Brazil; Yuguang Li, The University of Hong Kong, China; Ling Moi Lin, Sing Health, Singapore; Caline Mattar, World Health Professions Alliance, USA; MaryLouise McLaws, University of New South Wales, Australia; Geeta Mehta, Journal of Patient Safety and Infection Control, India; Shaheen Mehtar, Infection Control Africa Network, South Africa; Ziad Memish, Ministry of Health, Saudi Arabia; Babacar Ndoye, Infection Control Africa Network, Senegal; Fernando Otaiza, Ministry of Health, Chile; Diamantis Plachouras, European Centre for Disease Prevention and Control, Sweden; Maria Clara Padoveze, School of Nursing, University of São Paulo, Brazil; Mathias Pletz, Jena University, Germany; Marina Salvadori, Public Health Agency of Canada, Canada; Ingrid Schoeman, TB Proof, South Africa; Mitchell Schwaber, Ministry of Health, Israel; Nandini Shetty, Public Health England, United Kingdom; Mark Sobsey, University of North Carolina, USA; Paul Ananth Tambyah, National University Hospital, Singapore; Andreas Voss, Canisus-Wilhelmina Ziekenhuis, The Netherlands; Walter Zingg, University of Geneva Hospitals, Switzerland.
2. Experts from UNICEF:
Maya Ariei, Gregory Built, Simone Carter, Carlos Navarro Colorado, Anne Detjen, Nada Elattar, Maria Agnese Giordano, Gagan Gupta, Nagwa Hasanin, Linda Jones, Raoul Kamadjeu, Sarah Karmin, Asma Maladwala, Ana Nieto, Luwei Pearson Sarah Karmin, Jerome Pfaffmann.
3. External reviewers:
Susanna Esposito, World Association for Infectious Diseases and Immunological Disorders and University of Parma, Italy; Angela Dramowski, Stellenbosch University, South Africa; Alfredo Tagarro, Universidad Europea de Madrid, Spain.

Reviewers from IPA:

Berthold Koletzko, Ludwig-Maximilians-Universität München, Germany; Margaret Fisher, Department of Pediatrics, Monmouth Medical Center, USA; Jonathan Klein, University of Illinois at Chicago, IPA Executive Committee and Focal Point for WHO Collaboration, USA; Jane E Lucas, International Health and Child Development IPA Strategic Advisory Group on Early Child Development, USA; Mohamad Mikati, Division of Pediatric Neurology and Developmental Medicine, Duke University Medical Center, USA; Aman Pulungan, University of Indonesia, and Indonesian Pediatric Society and Asia Pacific Pediatric Association; Susan M Sawyer, International Association for Adolescent Health and

University of Melbourne, Australia; Mortada El-Shabrawi, Cairo University, Egypt; Russell M Viner, Faculty of Population Health Sciences of University College London and Royal College of Pediatrics and Child Health, UK.

4. WHO Secretariat:

Benedetta Allegranzi, Gertrude Avortri, Mekdim Ayana, Hanan Balkhy, April Baller, Elizabeth Barrera-Cancedda, Alessandro Cassini, Giorgio Cometto, Ana Paula Coutinho Rehse, Sophie Harriet Dennis, Sergey Eremin, Dennis Nathan Ford, Jonas Gonseth-Garcia, Rebecca Grant, Tom Grein, Ivan Ivanov, Landry Kabego, Pierre Claver Kariyo, Ying Ling Lin, Olivier Le Poulin, Ornella Lincetto, Abdi Rahman Mahamud, Madison Moon, Takeshi Nishijima, Pillar Ramon-Pardo Alice Simniceanu, Valeska Stempliuik, Maha Talaat Ismail, Joao Paulo Toledo, Anthony Twywan, Maria Van Kerkhove, Vicky Willet, Masahiro Zakoji, Bassim Zayed, Wilson Were.

References

1. World Health Organization. Advice on the use of masks in the context of COVID-19. Geneva: World Health Organization; 2020 (<https://apps.who.int/iris/handle/10665/331693> accessed 20 August 2020).
2. UNICEF. Convention on the Rights of the Child text. 1990 (<https://www.unicef.org/child-rights-convention/convention-text> accessed 20 August 2020).
3. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>.
4. Guan WJ, Ni ZY, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med*. 2020;382(18):1708-20. Epub 2020/02/29.
5. Wortham JM, Lee JT, Althomsons S, et al. Characteristics of Persons Who Died with COVID-19 - United States, February 12-May 18, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(28):923-9. Epub 2020/07/17.
6. European Centre for Disease Prevention and Control. COVID-19 in children and the role of school settings in COVID-19 transmission. 6 August 2020 (<https://www.ecdc.europa.eu/sites/default/files/documents/COVID-19-schools-transmission-August%202020.pdf> accessed 20 August 2020).
7. CDC COVID-19 Response Team. Coronavirus Disease 2019 in Children - United States, February 12-April 2, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(14):422-6. Epub 2020/04/10.
8. Ladhani SN, Amin-Chowdhury Z, Davies HG, et al. COVID-19 in children: analysis of the first pandemic peak in England. *Arch Dis Child*. 2020:archdischild-2020-320042.
9. Joint IPA-UNICEF COVID-19 Information Brief. Epidemiology, Spectrum, and Impact of COVID-19 on Children, Adolescents, and Pregnant Women. (<https://ipa-world.org/society-resources/code/images/HjNYEyfuM250.pdf> accessed 20 August 2020).
10. L'Huillier AG, Torriani G, Pigny F, Kaiser L, Eckerle I. Culture-Competent SARS-CoV-2 in Nasopharynx of Symptomatic Neonates, Children, and Adolescents. *Emerg Infect Dis*. 2020;26(10). Epub 2020/07/01.
11. Goldstein E, Lipsitch M, Cevik M. On the effect of age on the transmission of SARS-CoV-2 in households, schools and the community. *medRxiv*. 2020. (<https://www.medrxiv.org/content/10.1101/2020.07.19.20157362v2> accessed 20 August 2020).
12. Li X, Xu W, Dozier M, et al. The role of children in transmission of SARS-CoV-2: A rapid review. *J Glob Health*. 2020;10(1):011101. Epub 2020/07/03.
13. Ludvigsson JF. Children are unlikely to be the main drivers of the COVID-19 pandemic - A systematic review. *Acta Paediatr*. 2020;109(8):1525-30. Epub 2020/05/21.
14. Viner M, Mytton O, Bonnell C, et al. Susceptibility to and transmission of COVID-19 amongst children and adolescents compared with adults: a systematic review and meta-analysis. *medRxiv*. 2020. (<https://www.medrxiv.org/content/10.1101/2020.05.20.20108126v1> accessed 20 August 2020).
15. Macartney K, Quinn HE, Pillsbury AJ, Koirala A, Deng L, Winkler N, et al. Transmission of SARS-CoV-2 in Australian educational settings: a prospective cohort study. *Lancet Child Adolesc Health*. 2020. Epub 2020/08/08.
16. Fontanet A, Grant R, Tondeur L, et al. SARS-CoV-2 infection in primary schools in northern France: A retrospective cohort study in an area of high transmission. *medRxiv*. 2020. (<https://www.medrxiv.org/content/10.1101/2020.06.25.20140178v2> accessed 20 August 2020).
17. Fontanet A, Tondeur L, Madec Y et al. Cluster of COVID-19 in northern France: A retrospective closed cohort study. *medRxiv*. 2020. (<https://www.medrxiv.org/content/10.1101/2020.04.18.20071134v1> accessed 20 August 2020)
18. Stein-Zamir C, Abramson N, Shoob H, et al. A large COVID-19 outbreak in a high school 10 days after schools' reopening, Israel, May 2020. *Euro Surveill*. 2020;25(29). Epub 2020/07/29.
19. Torres JP, Pinera C, De La Maza V, et al. SARS-CoV-2 antibody prevalence in blood in a large school community subject to a Covid-19 outbreak: a cross-sectional study. *Clin Infect Dis*. 2020. Epub 2020/07/11.
20. Heavey L, Casey G, Kelly C, Kelly D, McDarby G. No evidence of secondary transmission of COVID-19 from children attending school in Ireland, 2020. *Euro Surveill*. 2020;25(21). Epub 2020/06/04.
21. Stringhini S, Wisniak A, Piumatti G, et al. Seroprevalence of anti-SARS-CoV-2 IgG antibodies in Geneva, Switzerland (SEROCoV-POP): a population-based study. *Lancet*. 2020;396(10247):313-9. Epub 2020/06/15.

22. Public Health England. Weekly Coronavirus Disease 2019 (COVID-19) Surveillance Report. Summary of COVID-19 surveillance systems. 2020.
23. Streeck H, Schulte B, Kümmerer B, et al. Infection fatality rate of SARS-CoV-2 infection in a German community with a super-spreading event. medRxiv. 2020 (<https://www.medrxiv.org/content/10.1101/2020.05.04.20090076v2> accessed 20 August 2020).
24. Shakiba M, Nazari S, Mehrabian F, et al. Seroprevalence of COVID-19 virus infection in Guilan province, Iran. medRxiv. 2020 (<https://www.medrxiv.org/content/10.1101/2020.04.26.20079244v1> accessed 20 August 2020).
25. Lu Y, Li Y, Deng W, et al. Symptomatic Infection is Associated with Prolonged Duration of Viral Shedding in Mild Coronavirus Disease 2019: A Retrospective Study of 110 Children in Wuhan. *Pediatr Infect Dis J.* 2020;39(7):e95-e9. Epub 2020/05/08.
26. Danis K, Epaulard O, Benet T, et al. Cluster of Coronavirus Disease 2019 (COVID-19) in the French Alps, February 2020. *Clin Infect Dis.* 2020;71(15):825-32. Epub 2020/04/12.
27. Xu Y, Li X, Zhu B, , et al. Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. *Nat Med.* 2020;26(4):502-5. Epub 2020/04/15.
28. Heald-Sargent T, Muller WJ, Zheng X, Rippe J, Patel AB, Kociolek LK. Age-Related Differences in Nasopharyngeal Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Levels in Patients With Mild to Moderate Coronavirus Disease 2019 (COVID-19). *JAMA Pediatr.* 2020. Epub 2020/08/04.
29. Jones TC, Mühlemann B, Veith T, et al. An analysis of SARS-CoV-2 viral load by patient age. medRxiv. 2020 (<http://medrxiv.org/lookup/doi/10.1101/2020.06.08.20125484> accessed 20 August 2020).
30. Canini L, Andreoletti L, Ferrari P, et al. Surgical mask to prevent influenza transmission in households: a cluster randomized trial. *PLoS One.* 2010;5(11):e13998. Epub 2010/11/26.
31. Chen X, Ran L, Liu Q, Hu Q, Du X, Tan X. Hand Hygiene, Mask-Wearing Behaviors and Its Associated Factors during the COVID-19 Epidemic: A Cross-Sectional Study among Primary School Students in Wuhan, China. *Int J Environ Res Public Health.* 2020;17(8). Epub 2020/04/26.
32. Simmerman JM, Suntaratiwong P, Levy J, et al. Findings from a household randomized controlled trial of hand washing and face masks to reduce influenza transmission in Bangkok, Thailand. *Influenza Other Respir Viruses.* 2011;5(4):256-67. Epub 2011/06/10.
33. Suess T, Remschmidt C, Schink SB, et al. The role of facemasks and hand hygiene in the prevention of influenza transmission in households: results from a cluster randomised trial; Berlin, Germany, 2009-2011. *BMC Infect Dis.* 2012;12:26. Epub 2012/01/28.
34. Uchida M, Kaneko M, Hidaka Y, et al. Effectiveness of vaccination and wearing masks on seasonal influenza in Matsumoto City, Japan, in the 2014/2015 season: An observational study among all elementary schoolchildren. *Prev Med Rep.* 2017;5:86-91. Epub 2016/12/17.
35. van der Sande M, Teunis P, Sabel R. Professional and home-made face masks reduce exposure to respiratory infections among the general population. *PLoS One.* 2008;3(7):e2618. Epub 2008/07/10.
36. Allison MA, Guest-Warnick G, Nelson D, et al. Feasibility of elementary school children's use of hand gel and facemasks during influenza season. *Influenza Other Respir Viruses.* 2010;4(4):223-9. Epub 2010/09/15.
37. Stebbins S, Downs JS, Vukotich CJ, Jr. Using nonpharmaceutical interventions to prevent influenza transmission in elementary school children: parent and teacher perspectives. *J Public Health Manag Pract.* 2009;15(2):112-7. Epub 2009/02/10.
38. Smart NR, Horwell CJ, Smart TS, Galea KS. Assessment of the Wearability of Facemasks against Air Pollution in Primary School-Aged Children in London. *Int J Environ Res Public Health.* 2020;17(11). Epub 2020/06/06.
39. Fikenzler S, Uhe T, Lavall D, et al. Effects of surgical and FFP2/N95 face masks on cardiopulmonary exercise capacity. *Clin Res Cardiol.* 2020. Epub 2020/07/08.
40. World Health Organization. Transmission of SARS-CoV-2: implications for infection prevention precautions. Geneva: World Health Organization; 2020 (<https://www.who.int/publications/i/item/modes-of-transmission-of-virus-causing-covid-19-implications-for-ipc-precaution-recommendations> accessed 20 August 2020).
41. Shelov S AT. *Caring for Your Baby and Young Child: Birth to Age 5*, Fifth Edition. American Academy of Pediatrics. Elk Grove Village, IL. . 2009.
42. Centers for Disease Control and Prevention. Considerations for Wearing Masks. United States of America; 2020 (<https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/cloth-face-cover-guidance.html> accessed 14 August 2020).
43. Swiss Federal Office for Public Health. New coronavirus: Masks. Koniz; 2020 (<https://www.bag.admin.ch/bag/en/home/krankheiten/ausbrueche-epidemien-pandemien/aktuelle-ausbrueche-epidemien/novel-cov/masken.html> accessed 20 August 2020).
44. Department of Health and Social Care. Face coverings: when to wear one and how to make your own. United Kingdom; 2020 (<https://www.gov.uk/government/publications/face-coverings-when-to-wear-one-and-how-to-make-your-own/face-coverings-when-to-wear-one-and-how-to-make-your-own> accessed 20 August 2020).
45. American Academy of Pediatrics. Cloth Face Coverings for Children During COVID-19. 2020 (<https://www.healthychildren.org/English/health-issues/conditions/COVID-19/Pages/Cloth-Face-Coverings-for-Children-During-COVID-19.aspx> accessed 20 August 2020).

46. Centers for Disease Control and Prevention. If You Are Immunocompromised, Protect Yourself From COVID-19. United States of America;2020 (<https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/immunocompromised.html> accessed 20 August 2020).
47. Cystic Fibrosis Foundation. COVID-19 Community Questions and Answers. 2020 (<https://www.cff.org/Life-With-CF/Daily-Life/Germs-and-Staying-Healthy/CF-and-Coronavirus/COVID-19-Community-Questions-and-Answers/#:~:text=People%20with%20CF%20should%20continue,cross%2Dinfection%20from%20CF%20germs.> accessed 20 August 2020).
48. Esposito S, Principi N. To mask or not to mask children to overcome COVID-19. Eur J Pediatr. 2020. Epub 2020/05/11.
49. Del Valle SY, Tellier R, Settles GS, Tang JW. Can we reduce the spread of influenza in schools with face masks? Am J Infect Control. 2010;38(9):676-7. Epub 2010/07/08.
50. American Cochlear Implant Alliance. Consideration of face shields as a return to school option. 2020 (<https://www.acialliance.org/page/consideration-of-face-shields-as-return-to-school-option> accessed 20 August 2020).
51. United Nations. Transparent masks aid communication for hard of hearing. 2020 (<https://www.un.org/en/coronavirus/transparent-masks-aid-communication-hard-hearing> accessed 20 August 2020).
52. Lindsley WG, Noti JD, Blachere FM, Szalajda JV, Beezhold DH. Efficacy of face shields against cough aerosol droplets from a cough simulator. J Occup Environ Hyg. 2014;11(8):509-18. Epub 2014/01/29.
53. Hirschmann MT, Hart A, Henckel J, Sadoghi P, Seil R, Mouton C. COVID-19 coronavirus: recommended personal protective equipment for the orthopaedic and trauma surgeon. Knee Surg Sports Traumatol Arthrosc. 2020;28(6):1690-8. Epub 2020/04/29.
54. Anon JB, Denne C, Rees D. Patient-Worn Enhanced Protection Face Shield for Flexible Endoscopy. Otolaryngol Head Neck Surg. 2020;163(2):280-3. Epub 2020/06/10.
55. Kähler CJ, Hain R. Fundamental protective mechanisms of face masks against droplet infections. Journal of Aerosol Science 148, 105617. (<https://doi.org/10.1016/j.jaerosci.2020.105617> accessed 20 August 2020).
56. World Health Organization. Rational use of personal protective equipment for coronavirus disease 2019 (COVID-19). Geneva: World Health Organization; 2020 (<https://apps.who.int/iris/rest/bitstreams/1274340/retrieve> accessed 20 August 2020).
57. Tony Blair Institute for Global Change. The Role of Face Shields in Responding to Covid-19. 2020 (<https://institute.global/sites/default/files/articles/The-Role-of-Face-Shields-in-Responding-to-Covid-19.pdf> accessed 20 August, 2020).
58. Perencevich EN, Diekema DJ, Edmond MB. Moving Personal Protective Equipment Into the Community: Face Shields and Containment of COVID-19. JAMA. 2020. Epub 2020/04/30.
59. Victoria State Health and Human Services. Face coverings: whole of Victoria. 2020 (<https://www.dhhs.vic.gov.au/face-coverings-covid-19#what-does-wearing-a-face-covering-mean> accessed 20 August 2020).
60. Ministry of Health. Guidance for use of masks and face shields. Singapore;2020 (<https://www.moh.gov.sg/news-highlights/details/guidance-for-use-of-masks-and-face-shields> accessed 20 August 2020).
61. Roberge RJ. Face shields for infection control: A review. J Occup Environ Hyg. 2016;13(4):235-42. Epub 2015/11/13.
62. World Health Organization. Disease Commodity Package v5. Geneva: World Health Organization; 2020 (<https://www.who.int/emergencies/what-we-do/prevention-readiness/disease-commodity-packages/dcp-ncov.pdf?ua=1> accessed August 20, 2020).

WHO and UNICEF continue to monitor the situation closely for any changes that may affect this interim guidance. Should any factors change, an update will be issued. Otherwise, this interim guidance document will expire 2 years after the date of publication.

© World Health Organization and the United Nations Children’s Fund (UNICEF), 2020. Some rights reserved. This work is available under the [CC BY-NC-SA 3.0 IGO](https://creativecommons.org/licenses/by-nc-sa/3.0/) licence.

WHO reference number: [WHO/2019-nCoV/IPC_Masks/Children/2020.1](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/operational-recommendations/WHO-2019-nCoV-IPC_Masks/Children/2020.1)